

LETTER • OPEN ACCESS

## Increased importance of methane reduction for a 1.5 degree target

To cite this article: William J Collins *et al* 2018 *Environ. Res. Lett.* **13** 054003

View the [article online](#) for updates and enhancements.

### Related content

- [Impact of short-lived non-CO<sub>2</sub> mitigation on carbon budgets for stabilizing global warming](#)  
Joeri Rogelj, Malte Meinshausen, Michiel Schaeffer *et al.*
- [Drivers and patterns of land biosphere carbon balance reversal](#)  
Christoph Müller, Elke Stehfest, Jelle G van Minnen *et al.*
- [Sensitivity of carbon budgets to permafrost carbon feedbacks and non-CO<sub>2</sub> forcings](#)  
Andrew H MacDougall, Kirsten Zickfeld, Reto Knutti *et al.*

### Recent citations

- [Methane emissions from on-road vehicles in China: a case study in an urban tunnel](#)  
Yanli Zhang *et al*
- [Demonstrating GWP\\*: a means of reporting warming-equivalent emissions that captures the contrasting impacts of short- and long-lived climate pollutants](#)  
John Lynch *et al*
- [Stable climate metrics for emissions of short and long-lived species—combining steps and pulses](#)  
William J Collins *et al*

# Environmental Research Letters



## LETTER

### OPEN ACCESS

RECEIVED  
30 January 2018

REVISED  
9 March 2018

ACCEPTED FOR PUBLICATION  
22 March 2018

PUBLISHED  
20 April 2018

Original content from  
this work may be used  
under the terms of the  
[Creative Commons  
Attribution 3.0 licence](#).

Any further distribution  
of this work must  
maintain attribution to  
the author(s) and the  
title of the work, journal  
citation and DOI.



## Increased importance of methane reduction for a 1.5 degree target

William J Collins<sup>1,6</sup> , Christopher P Webber<sup>1</sup>, Peter M Cox<sup>2</sup>, Chris Huntingford<sup>3</sup>, Jason Lowe<sup>4,5</sup>, Stephen Sitch<sup>2</sup>, Sarah E Chadburn<sup>2,5</sup>, Edward Comyn-Platt<sup>3</sup>, Anna B Harper<sup>2</sup>, Garry Hayman<sup>3</sup> and Tom Powell<sup>2</sup>

<sup>1</sup> University of Reading, Reading, RG6 6BB, United Kingdom

<sup>2</sup> University of Exeter, Exeter EX4 4QF, United Kingdom

<sup>3</sup> Centre for Ecology and Hydrology, Wallingford, Oxfordshire, OX10 8BB, United Kingdom

<sup>4</sup> Met Office Hadley Centre, Fitzroy Road, Exeter EX1 3PB, United Kingdom

<sup>5</sup> University of Leeds, Leeds LS2 9JT, United Kingdom

<sup>6</sup> Author to whom any correspondence should be addressed.

E-mail: [w.collins@reading.ac.uk](mailto:w.collins@reading.ac.uk)

**Keywords:** carbon budgets, methane mitigation, non-CO<sub>2</sub> greenhouse gases, climate targets

Supplementary material for this article is available [online](#)

### Abstract

To understand the importance of methane on the levels of carbon emission reductions required to achieve temperature goals, a process-based approach is necessary rather than reliance on the transient climate response to emissions. We show that plausible levels of methane (CH<sub>4</sub>) mitigation can make a substantial difference to the feasibility of achieving the Paris climate targets through increasing the allowable carbon emissions. This benefit is enhanced by the indirect effects of CH<sub>4</sub> on ozone (O<sub>3</sub>). Here the differing effects of CH<sub>4</sub> and CO<sub>2</sub> on land carbon storage, including the effects of surface O<sub>3</sub>, lead to an additional increase in the allowable carbon emissions with CH<sub>4</sub> mitigation. We find a simple robust relationship between the change in the 2100 CH<sub>4</sub> concentration and the extra allowable cumulative carbon emissions between now and 2100 ( $0.27 \pm 0.05$  GtC per ppb CH<sub>4</sub>). This relationship is independent of modelled climate sensitivity and precise temperature target, although later mitigation of CH<sub>4</sub> reduces its value and thus methane reduction effectiveness. Up to 12% of this increase in allowable emissions is due to the effect of surface ozone. We conclude early mitigation of CH<sub>4</sub> emissions would significantly increase the feasibility of stabilising global warming below 1.5 °C, alongside having co-benefits for human and ecosystem health.

### 1. Introduction

Meeting the Paris temperature targets by reducing CO<sub>2</sub> emissions alone represents a huge challenge, even for the more optimistic assessments of the allowable carbon budgets (Millar *et al* 2017). Most existing scenarios that avoid 2 °C of global warming, and almost all of those that avoid 1.5 °C, assume periods of negative global CO<sub>2</sub> emissions in order to stay within the implied cumulative carbon budgets (Rogelj *et al* 2015a). This is via the widespread deployment of carbon dioxide removal (CDR) (Smith 2016) which might not be as effective as assumed (Harper *et al* 2018). Any additional options for mitigating greenhouse gases can therefore increase the feasibility of this challenge.

The transient climate response to emissions (TCRE) has proved useful in illustrating the dependence of temperature on the cumulative emissions of CO<sub>2</sub>. However care needs to be taken as the scenarios used in the IPCC 5th Assessment Report (AR5) (Pachauri *et al* 2014) assumed specific changes in non-CO<sub>2</sub> agents such as aerosols and CH<sub>4</sub>. These calculations also did not include biogeochemical feedbacks that might affect the concentrations of the greenhouse gases such as changes in permafrost and wetlands (Comyn-Platt *et al* 2018). The relationship between cumulative carbon emissions and global temperature target will therefore depend crucially on the future mix of CO<sub>2</sub> and non-CO<sub>2</sub> agents which may differ significantly from that assumed in AR5. As a consequence cumulative carbon budgets are very

sensitive to assumptions in scenarios for non-CO<sub>2</sub> greenhouse gases (Rogelj *et al* 2015b).

Mitigation of anthropogenic CH<sub>4</sub> emissions leads to rapid decreases in its concentration, with an approximately 12 year response time. CH<sub>4</sub> mitigation therefore offers potential for rapidly reducing climate warming, either in the near-term to prevent a temporary exceedance of the 1.5 or 2.0 °C peak warming threshold, or later in the century to bring down temperatures after an overshoot of temperature to higher levels. A recent study (Stohl *et al* 2015) found that inexpensive or even cost negative CH<sub>4</sub> mitigation options could reduce 2050 temperatures by 0.25 °C.

Methane has a direct radiative forcing of climate. It is the second largest contributor to anthropogenic forcing over the historical period, and its atmospheric chemistry leads to O<sub>3</sub> and water vapour, themselves GHGs, adding to the forcing (Myhre *et al* 2013). Changes to atmospheric CH<sub>4</sub>, O<sub>3</sub> and CO<sub>2</sub> will also affect the ocean and land carbon cycles, through direct warming effects (climate-carbon feedbacks), increasing the rates of plant respiration and decomposition of soil organic carbon. There are also indirect physiological effects of O<sub>3</sub>, decreasing, and CO<sub>2</sub>, increasing, plant productivity and hence carbon uptake (Sitch *et al* 2007, Collins *et al* 2010, Sitch *et al* 2008). These carbon-cycle effects are typically included in calculations of the effects of CO<sub>2</sub> emissions, but are currently ignored when calculating the CO<sub>2</sub>-equivalence of non-CO<sub>2</sub> gases such as CH<sub>4</sub> (MacDougall *et al* 2013). Recent studies (Collins *et al* 2013, Gasser *et al* 2017) estimated that the climate-carbon cycle feedbacks increase the temperature impacts of CH<sub>4</sub> by around 20% on 100 year timescales.

As a result of these typically-neglected effects, it has been argued that the total carbon budget for stabilization of the climate at about 2 °C might be much more sensitive to the atmospheric concentration of CH<sub>4</sub> than hereto expected (Cox and Jeffery 2010). This is likely to be even more so for a 1.5 °C target. This is because the impact on land carbon storage arising from a change in radiative forcing due to mitigation of CO<sub>2</sub> differs significantly from the impact of a similar non-CO<sub>2</sub> radiative forcing mitigation (Huntingford *et al* 2011). When including the damaging effects of surface O<sub>3</sub>, reductions in the emissions of CH<sub>4</sub> have the potential to significantly increase land carbon storage.

## 2. Methods

### 2.1. IMOGEN-JULES


To understand the potential additional benefits of CH<sub>4</sub> reductions on allowable cumulative carbon emissions consistent with the Paris targets, we use the Joint UK Land-Environment Simulator (JULES) (Clark *et al* 2011) coupled with the intermediate complexity climate model IMOGEN ‘Integrated Model Of Global Effects of climatic aNomalies’ (Huntingford *et al* 2010).

The combined IMOGEN-JULES framework thus provides an intermediate complexity climate-carbon modelling system. IMOGEN utilises ‘pattern-scaling’ to capture the main features of expected local and monthly meteorological changes interpolated to alternative future levels of global warming. This is connected to a gridded version of the land surface model JULES (version 4.8) (Clark *et al* 2011) to understand the impacts of any transition to different stable warming levels.

IMOGEN comprises a global energy balance model (EBM) whose global climate response characteristics (climate sensitivity for land and ocean, ocean diffusivity etc.) can be chosen to represent any global climate model (GCM). It is driven by time-series of CO<sub>2</sub> concentrations and non-CO<sub>2</sub> radiative forcing. IMOGEN generates gridded outputs of monthly anomaly fields of surface temperature, precipitation, humidity, wind-speed, surface shortwave and longwave radiation and pressure. These anomalies are derived by scaling the patterns from the output from each GCM, assuming these are linear in global surface temperature change. Here the data from the 34 CMIP5 GCMs running the RCP8.5 scenario (Taylor *et al* 2013) are used to derive both the global climate characteristics and climate patterns. Although the greenhouse gas forcings used in this study will be closer to the RCP2.6 scenario, the RCP8.5 scenario was used to get the clearest signal to determine the climate patterns.

The JULES configuration also includes modelled O<sub>3</sub> damage to photosynthesis, affecting land-atmosphere CO<sub>2</sub> exchange (Sitch *et al* 2007). This O<sub>3</sub> damage parameterisation can be set to ‘low’ or ‘high’ sensitivity to span the uncertainty in our knowledge of the sensitivity of plants globally. We also include a ‘no’ sensitivity to allow the separation of the ozone effect. In this study we use the low sensitivity parameterisation as the standard configuration, with separate tests of the effects of using the ‘no’ and ‘high’ sensitivities. Surface O<sub>3</sub> concentrations are parameterised as two-dimensional fields as a function of the global average CH<sub>4</sub> concentration. These are previously derived from global chemistry-climate simulations using the HadGEM3 model for global mean atmospheric CH<sub>4</sub> mixing ratios of 1285 ppb and 2062 ppb (Stohl *et al* 2015). Within IMOGEN-JULES, the O<sub>3</sub> concentration is calculated at each grid point as a function of CH<sub>4</sub> using a linear interpolation between O<sub>3</sub> concentrations at the above mixing ratios.

To set the initial (2015) conditions for the land carbon stores, the IMOGEN-JULES model is spun up for 1000 years at 1850 conditions and then run to 2015 with prescribed historical CO<sub>2</sub> mixing ratios, land use, and global surface temperatures from Morice *et al* (2012) (reaching 0.89 °C by 2015). The spin up and historical simulation are carried out for each climate model realisation. For this study we invert the IMOGEN-JULES configuration, running forward from 2015 with specified global temperature profiles,



and specified non- $\text{CO}_2$  radiative forcing changes from 2015. IMOGEN-JULES derives the  $\text{CO}_2$  concentrations in each year from the EBM calculations and thence the uptake by the land biosphere; the global carbon cycle is closed with a simple description of global oceanic draw-down of  $\text{CO}_2$  (Joos *et al* 1996). A control simulation is also run maintaining 1850 forcings and temperatures until 2100. Further details of the IMOGEN-JULES setup and the inversion procedure can be found in Comyn-Platt *et al* (2018).

## 2.2. Temperature and methane scenarios

We determine the carbon budgets consistent with three specified temperature trajectories that stabilise at 1.5 $^{\circ}\text{C}$  (with and without overshoot) and 2.0 $^{\circ}\text{C}$  above

pre-industrial levels as shown in figure 1(a). These profiles are generated according to the algorithm in Huntingford *et al* (2017) as in Comyn-Platt *et al* (2018). The results are found not to be sensitive to the exact form of the temperature trajectories.

The future non- $\text{CO}_2$ , non- $\text{CH}_4$  radiative forcings are taken from one of the Shared Socio-economic Pathways (SSPs) SSP2–2.6 (O'Neill *et al* 2017, Riahi *et al* 2017) by subtracting the  $\text{CO}_2$  and  $\text{CH}_4$  (and associated  $\text{O}_3$  and stratospheric water vapour) contributions from the total SSP2–2.6 radiative forcing. We follow the prescription of these terms in the MAGICC climate model (Meinshausen *et al* 2011). After 2015, land-use is fixed at 2015 levels. Here, the IMOGEN physical parameters are varied to represent the climate

characteristics, such as the different climate sensitivities, of 34 CMIP5 models.

There is a wide range in the CH<sub>4</sub> emissions in the SSPs that achieve a forcing of 2.6 W m<sup>-2</sup> in 2100, suggesting that the options for mitigation are not exhausted (Gernaat *et al* 2015). We construct four different anthropogenic CH<sub>4</sub> mitigation scenarios (figure 1 (b)). The first three are ‘High’ CH<sub>4</sub> and ‘Medium’ CH<sub>4</sub> which span the highest and lowest of the SSP2–2.6, and ‘Low’ CH<sub>4</sub> which we parameterise as following the Medium scenario to 2020 then decaying faster to 62 Tg CH<sub>4</sub> yr<sup>-1</sup> by 2100. For the High CH<sub>4</sub> scenario, CH<sub>4</sub> concentrations increase following the an upper bound of SSP4–2.6 and SSP5–2.6 CH<sub>4</sub> concentration projections from the GCAM integrated assessment model (IAM) (Calvin *et al* 2017). For the Medium CH<sub>4</sub> scenario, concentrations follow SSP2–2.6 as generated by the IMAGE 3.0 IAM (van Vuuren *et al* 2017). For the Low CH<sub>4</sub> scenario, we assume extra reductions are possible by removing the restriction on cost minimisation. To generate a smooth curve we parameterise emissions (in Tg CH<sub>4</sub> yr<sup>-1</sup>) as  $55 + \frac{337.25}{x^{1.337}}$ , where  $x$  is the number of years after 2020. This projects a lower CH<sub>4</sub> projection curve than the strongest mitigation SSP storyline (SSP1–2.6 variants). The High, Medium and Low scenarios lead to year 2100 atmospheric CH<sub>4</sub> concentrations of 1839, 1275 and 1008 ppb, respectively. We also consider a fourth scenario ‘Late’, to test whether the timing of the CH<sub>4</sub> mitigation matters, where emissions are maintained at current (2015) levels until 2050 and then apply the same rate of mitigation for the Low CH<sub>4</sub> profile post-2015, but extended to ensure that the 2100 concentration matches Low CH<sub>4</sub>. Note that we are not assuming specific methane mitigation measures in these scenarios, or possible effects on co-emitted species such as N<sub>2</sub>O.

Emissions are converted into concentrations using the formulation of the MAGICC model (which includes natural emissions of 250 Tg CH<sub>4</sub> yr<sup>-1</sup>). Radiative forcings for the CH<sub>4</sub> scenarios are calculated using formulae including the short-wave absorption (Etminan *et al* 2016), and the overlap with N<sub>2</sub>O using the N<sub>2</sub>O concentrations in SSP2–2.6. The contributions from O<sub>3</sub> and stratospheric water vapour are added in as linear functions of CH<sub>4</sub> mixing ratio. From IPCC AR5 (Myhre *et al* 2013) these amount to  $2.36 \times 10^{-4} \pm 1.09 \times 10^{-4}$  Wm<sup>-2</sup> per ppb CH<sub>4</sub> (0.65 ± 0.3 times the CH<sub>4</sub> radiative efficiency).

This spread in possible CH<sub>4</sub> trajectories is wider than typically projected in integrated assessment models (IAMs) (Rogelj *et al* 2015a). However, the IAM outputs are unlikely to span the full range of CH<sub>4</sub> measures that are available. This is partly due to their cost minimisation approaches which exclude the more expensive measures and neglect the social costs of methane (Shindell *et al* 2017), and their lack of diversity in treatment of non-CO<sub>2</sub> mitigation measures. These IAMs also have limited representation of the specific processes responsible for methane production and of

the technologies available for methane mitigation. It is therefore difficult to estimate how deep (or not) reductions can go. Achieving our most stringent scenario would be expected to draw on specific sectoral measures to address CH<sub>4</sub>. These could include increasing agricultural efficiency, decreased food waste and decreased beef consumption (van Vuuren *et al* 2017). The Low and Late scenarios should therefore be seen as illustrative examples.


### 3. Results

#### 3.1. Carbon budgets

For the High CH<sub>4</sub> scenario (no CH<sub>4</sub> mitigation) the allowable carbon emissions from 2015–2100 span from  $149 \pm 51$  GtC for 1.5 °C (no overshoot),  $143 \pm 56$  GtC for 1.5° with overshoot, to  $403 \pm 94$  GtC for the 2° temperature pathway. The uncertainty is due to the range of climate sensitivities of the CMIP5 models emulated by the IMOGEN framework. Rather than these absolute budgets we focus on the differences in the cumulative carbon emissions from the inversions for the different CH<sub>4</sub> scenarios. These show almost no dependence on the climate model realisation and little dependence on the temperature profile. The benefit of the Medium vs the High CH<sub>4</sub> scenario is approximately 155 GtC over the period 2015–2100 (figure 2(a)). Stronger CH<sub>4</sub> mitigation down to the Low scenario gains another 80 GtC if it is done early. The loss in benefit from delaying CH<sub>4</sub> mitigation according to the Late CH<sub>4</sub> scenario is 40 GtC. These values are similar to a study comparing no mitigation with stringent mitigation (Rogelj *et al* 2015b) which calculated an increase of 130 GtC in the carbon budget, with a 30 GtC penalty for late mitigation.

The relationship between the allowable carbon emissions from 2015–2100 and CH<sub>4</sub> concentrations at 2100 is almost linear (excluding the Late CH<sub>4</sub> scenario) with very little difference between the climate model realisations (figure 2(b)). The slopes are  $-0.269 \pm 0.001$  GtC ppb<sup>-1</sup> for the 1.5° and 1.5° overshoot profile and  $-0.277 \pm 0.002$  GtC ppb<sup>-1</sup> for the 2° C profile. Compared to the CH<sub>4</sub> forcing at 2100 (including the O<sub>3</sub> and stratospheric water vapour effects), this is equivalent to 350 or 360 GtC (Wm<sup>-2</sup>)<sup>-1</sup>. There are uncertainties in these relationships due to the uncertainty in the total radiative efficiency of methane. As these relationships are based on the methane concentrations, rather than emissions, uncertainties in the methane lifetime do not affect the result. The uncertainty in the direct methane radiative efficiency is taken to be 9% of the total (Etminan *et al* 2016). When combined with the 16% uncertainty from the ozone and water vapour contributions this leads to an overall uncertainty of 18%, (0.048 GtC ppb<sup>-1</sup>). This uncertainty includes within its span the relationship ( $-0.236$  GtC ppb<sup>-1</sup>) expected using the Myhre *et al* (1998) forcing instead of Etminan *et al* (2016).






**Figure 2.** Impact of CH<sub>4</sub> mitigation on the carbon budget for the three temperature profiles. (a) Increase in allowable carbon emissions compared to the High CH<sub>4</sub> scenario. Data are shown for the three temperature profiles. The widths of the lines cover the range of the CMIP5 models. (b) Difference in allowable carbon emissions between pairs of CH<sub>4</sub> scenarios, as a function of difference in CH<sub>4</sub> concentration for each year 2015–2100. The widths of the lines cover the range of the CMIP5 models. The dashed lines connect the differences in 2100 carbon budget against 2100 CH<sub>4</sub> concentrations for the Low, Medium and High CH<sub>4</sub> scenarios. For the Late vs High CH<sub>4</sub> scenario only the 1.5° temperature profile is shown.

The change in carbon budgets (high methane vs low methane) can be broken down into the different carbon stores: atmosphere, land (soil and vegetation) and ocean (figure 3(a)). We define the airborne fraction  $\alpha_F = \Delta\text{CO}_2 / \Delta E_{\text{CO}_2}$ , where  $\Delta\text{CO}_2$  is the change in the atmospheric CO<sub>2</sub> burden and  $\Delta E_{\text{CO}_2}$  is the change in cumulative CO<sub>2</sub> emissions, both in GtC. We find that the  $\alpha_F$  of the extra carbon allowed through CH<sub>4</sub> mitigation is independent of the climate sensitivity of each climate model.  $\alpha_F$  is also the same when comparing Low-High and Medium-High CH<sub>4</sub> mitigation (not shown). There is a slight dependence of  $\alpha_F$  on temperature profile with the 1.5°C profiles having an  $\alpha_F$  of 0.44 vs 0.49 for the 2°C profile. The Late CH<sub>4</sub>

mitigation does not follow the same linear relationship as the Low or Medium scenarios, falling well below the line of proportionality in figure 2(b). With late CH<sub>4</sub> mitigation, the comparative increase in allowable atmospheric CO<sub>2</sub> concentrations (compared to High CH<sub>4</sub>) does not occur until late in the century. The increase in the atmospheric carbon is the same as for the early mitigation, but the ocean and the land have not had time to take up this extra carbon and the  $\alpha_F$  of the extra CO<sub>2</sub> is thus higher (0.53).

Since surface O<sub>3</sub> decreases vegetation productivity, mitigation of CH<sub>4</sub> leads to additional climate benefits than might be expected simply through the radiative forcing. Decreasing atmospheric CH<sub>4</sub> concentrations



**Figure 3.** Difference in carbon stores in the atmosphere, ocean and land at 2100 compared to the High CH<sub>4</sub> scenario. (a) Low CH<sub>4</sub> scenario for the three temperature profiles, and the Late CH<sub>4</sub> scenario for the 1.5° temperature profile. Values shown are percentages of the total carbon stores (equal to allowable carbon emissions). Error bars are very small and show the inter-model standard deviation. (b) As (a), but for high O<sub>3</sub> sensitivity, showing the contributions of low and high O<sub>3</sub> sensitivity to the increased soil carbon. Diagonal hatch is low damage, total hatch is high damage, cross hatch is extra effect of high vs low damage.

reduces O<sub>3</sub> levels and increases the uptake of carbon into vegetation and soils. In terms of equation (1), reducing O<sub>3</sub> reduces  $\alpha_F$ . We test this through further inversions assuming no and high sensitivity of vegetation to O<sub>3</sub>, compared with the baseline parameterisation in the previous results of lower plant-O<sub>3</sub> sensitivity. We find that by increasing the impacts on the land carbon uptake, O<sub>3</sub> damage adds 9–28 GtC (4%–12%) to the benefit of the Low vs High CH<sub>4</sub> scenarios depending on the assumed sensitivity of vegetation to O<sub>3</sub> (figure 3(b)).

### 3.2. Linearity of carbon budgets

To maintain the radiative balance in the inverse model the change in atmospheric CO<sub>2</sub> is entirely determined by the change in the non-CO<sub>2</sub> forcing. Since we invert IMOGEN to derive the radiation balance consistent with the specified temperature profiles, the greenhouse gas forcing must be the same at any given time, such as at 2100, (assuming the climate sensitivities to radiative forcing from CH<sub>4</sub> and CO<sub>2</sub> are equal). So

$$\Delta F_{\text{CO}_2} + \Delta F_{\text{CH}_4} = 0, \text{ or } \Delta \text{CO}_2 \times A_{\text{CO}_2} + \Delta \text{CH}_4 \times \bar{A}_{\text{CH}_4} = 0;$$

where  $\Delta \text{CO}_2$  and  $\Delta \text{CH}_4$  are the CO<sub>2</sub> and CH<sub>4</sub> burdens in GtC and GtCH<sub>4</sub>, and  $\bar{A}_{\text{CH}_4}$  and  $\bar{A}_{\text{CO}_2}$  are the average radiative efficiencies for increases in CH<sub>4</sub> (including its indirect effects) and CO<sub>2</sub> in Wm<sup>-2</sup> GtCH<sub>4</sub><sup>-1</sup> or Wm<sup>-2</sup> GtC<sup>-1</sup>. So combining these with the airborne fraction  $\alpha_F$  defined previously gives the ratio of extra cumulative carbon emissions ( $\Delta E_{\text{CO}_2}$ ) to change in CH<sub>4</sub> abundance:


$$\frac{\Delta E_{\text{CO}_2}}{\Delta \text{CH}_4} = -\frac{\bar{A}_{\text{CH}_4}}{\alpha_F \bar{A}_{\text{CO}_2}} \quad (1)$$

This equation is exact and simply follows from the way we have defined  $\bar{A}_{\text{CH}_4}$ ,  $\bar{A}_{\text{CO}_2}$  and  $\alpha_F$ . The linear relationship between the change in the allowable emissions and the change in 2100 forcing therefore implies a constant ratio between the cumulative emissions to 2100 and the 2100 atmospheric CO<sub>2</sub> burden, i.e. a constant airborne fraction for the extra allowable emissions as found in figure 3(a). Although  $\bar{A}_{\text{CH}_4}$  and  $\bar{A}_{\text{CO}_2}$  are not constant, but functions of the atmospheric CO<sub>2</sub> levels and the magnitudes of the changes  $\Delta \text{CO}_2$  and  $\Delta \text{CH}_4$ , the deviations from linearity are small for the methane mitigation scenarios used here. The slightly higher  $\alpha_F$  for the 2.0° temperature profile is due to the lower radiative efficiency ( $\bar{A}_{\text{CO}_2}$ ) at higher absolute CO<sub>2</sub> levels.

The equation also holds in the more realistic case where the extra allowed CO<sub>2</sub> is not emitted with the time profile required to precisely follow the prescribed temperature curve, although in this case the  $\alpha_F$  may be slightly different from found in this study. The energy balance has little dependence on the shape of the temperature curve before 2100 (or any specific time), and is dominated by the absolute temperature and its time derivative at 2100. This relationship has no dependence on climate sensitivity. However the  $\alpha_F$  will be affected by the sensitivity of the carbon cycle to changes in atmospheric CO<sub>2</sub>, surface temperature and precipitation (Arora *et al* 2013).

Allen *et al* (2016) have derived a variant of the Global Warming Potential metric (GWP\*) that relates the change in *cumulative* emissions of CO<sub>2</sub> to the change in *instantaneous* emissions of a short-lived species (here CH<sub>4</sub>).

$$\text{GWP}^* = \frac{\Delta E_{\text{CO}_2}}{\Delta e_{\text{CH}_4}} = \frac{\text{AGWP}_{\text{CH}_4}^{\text{H}}}{\text{AGWP}_{\text{CO}_2}^{\text{H}}/H},$$



**Figure 4.** (a) Effect of CH<sub>4</sub> mitigation (Low-High) on surface ozone levels. (b) Effect of CH<sub>4</sub> mitigation (Low-High) on global NPP where vegetation has no sensitivity to O<sub>3</sub>, low sensitivity (as standard setup) and high sensitivity to ozone. The widths of the lines cover the range of the CMIP5 models. (c) Map of the regions of increased NPP attributable to reducing surface O<sub>3</sub> (Low CH<sub>4</sub> vs High CH<sub>4</sub> scenarios) using the high sensitivity to O<sub>3</sub>, as a percentage of the total NPP.

where  $\Delta e_{\text{CH}_4}$  is the change in the instantaneous CH<sub>4</sub> emission rate (in GtCH<sub>4</sub> yr<sup>-1</sup>),  $H$  is a chosen time-frame, and AGWP<sub>X</sub><sup>H</sup> are the absolute GWPs for CH<sub>4</sub> and CO<sub>2</sub>. The absolute GWPs can be expanded to give:

$$\frac{\Delta E_{\text{CO}_2}}{\Delta e_{\text{CH}_4}} = \frac{\overline{A}_{\text{CH}_4} \tau_{\text{CH}_4} (1 - e^{-H/\tau_{\text{CH}_4}})}{\overline{A}_{\text{CO}_2} \times \alpha_F(H)}$$



where  $\alpha_F(H)$  is the airborne fraction of a pulse of  $\text{CO}_2$  averaged over  $H$  years. This is similar to equation (1), but only equal to it if the  $\text{CH}_4$  has reached equilibrium (i.e.  $\Delta\text{CH}_4$  can be replaced by  $\Delta e_{\text{CH}_4} \times \tau_{\text{CH}_4}$ ) and the airborne fraction of  $\text{CO}_2$  in the  $\text{AGWP}_{\text{CO}_2}^H$  ( $\alpha_F(H)$ ) is equal to the  $\alpha_F$  of the extra allowed  $\text{CO}_2$ .

In terms of  $\text{GWP}^*$ , the results from our experiments give a ratio  $\frac{\Delta E_{\text{CO}_2}}{\Delta e_{\text{CH}_4}}$  at the end of the century of 2900 (low-medium mitigation) to 3300 (low-high mitigation) in  $\text{GtCO}_2 \text{ GtCH}_4^{-1} \text{ yr}^{-1}$ , which compares well with a  $\text{GWP}^*$  (100 years) of 2800 yr, given that implicit in the  $\text{GWP}^*$  approximation are the assumptions that the  $\text{CH}_4$  concentrations have equilibrated and that the  $\text{CO}_2$  airborne fraction is constant.

### 3.3. Air quality and productivity benefits

We find that  $\text{CH}_4$  mitigation has non-climate benefits in terms of air quality and vegetation productivity (by allowing greater atmospheric  $\text{CO}_2$  levels, and by reducing the damage from  $\text{O}_3$ ). West *et al* (2012) found that a strong methane mitigation scenario (emission decrease of  $180 \text{ Tg CH}_4 \text{ yr}^{-1}$ ) resulted in a decrease in global ozone concentrations of around 2 ppb and avoided mortalities of around 90 000 per year. In this study, mitigation by  $260 \text{ TgCH}_4 \text{ yr}^{-1}$  (Low vs High scenario) achieves a decrease in surface  $\text{O}_3$  concentration of 3 ppb as a global average, with the largest impact in the tropics (see figure 4(a)). Therefore a rough scaling of West *et al* (2012) would suggest a benefit of around 130 000 avoided mortalities per year.

The increased allowable  $\text{CO}_2$  levels lead to increased net primary plant productivity (NPP) in JULES by 4% as a global average (figure 4(b)). If we assume the high sensitivity of plants to ozone the effects of  $\text{O}_3$  reduction add up to another 2% increase in NPP globally. In places where the changes in ozone overlap with areas of high productivity (Eastern US, northern Europe) the reductions in ozone could increase total NPP by 4%–6% in the high sensitivity case (figure 4(c)).

## 4. Conclusions

We conclude that mitigating  $\text{CH}_4$  can lead to substantial benefits in the allowable carbon emissions consistent with either a  $1.5^\circ$  or  $2.0^\circ$  temperature target. We find a robust relationship between decreased  $\text{CH}_4$  concentrations at the end of the century and increased budget of allowable carbon emissions to 2100. This relationship is independent of climate sensitivity or temperature pathway. These changes come from the direct radiative effects of  $\text{CH}_4$  and its atmospheric oxidation products, from the carbon uptake by the land and ocean, and from the effects of  $\text{O}_3$  on plant productivity. Budget calculations based simply on TCRE will therefore underestimate allowed emissions. As well as making carbon targets more feasible,  $\text{CH}_4$

mitigation leads to substantial land ecosystem benefits through increased productivity, and to improved air quality. The variation in  $\text{CH}_4$  emissions between the IAMs in the SSP scenarios shows that there is substantial opportunity for  $\text{CH}_4$  mitigation even using the cost optimisation assumptions in these models. Very large cuts in  $\text{CO}_2$  emissions will certainly be needed to achieve the climate goals, but our study shows that the benefits of  $\text{CH}_4$  mitigation could be substantially larger than the IAMs assume, making the exploration and costing of more ambitious reduction potentials and their co-benefits a priority.

## Acknowledgments

The work was undertaken as part of the UK Natural Environment Research Council's programme 'Understanding the Pathways to and Impacts of a  $1.5^\circ\text{C}$  Rise in Global Temperature' through grants NE/P014909/1, MOC1.5 (WC, CW, CH, PC, SS, JL), NE/P015050/1 CLIFFTOP (EC-P, GH, SC), and NE/P014941/1 CLUES (PC, TP). AH acknowledges support from the EPSRC Fellowship 'Negative Emissions and the Food-Energy-Water Nexus' (EP/N030141/1). WC also acknowledges support under Research Council of Norway, project no. 235548.

## ORCID iDs

William J Collins  <https://orcid.org/0000-0002-7419-0850>

## References

- Allen M R, Fuglestedt J S, Shine K P, Reisinger A, Pierrehumbert P T and Forster P M 2016 New use of global warming potentials to compare cumulative and short-lived climate pollutants *Nat. Clim. Change* **6** 773–6
- Arora V K *et al* 2013 Carbon-concentration and carbon-climate feedbacks in CMIP5 Earth system models *J. Clim.* **26** 5289–314
- Calvin K *et al* 2017 The SSP4: a world of deepening inequality *Glob. Environ. Change* **42** 284–96
- Clark D *et al* 2011 The Joint UK Land Environment Simulator (JULES), model description - Part 2: carbon fluxes and vegetation dynamics *Geosci. Model Dev.* **4** 701–22
- Collins W J, Fry M M, Yu H, Fuglestedt J S, Shindell D T and West J J 2013 Global and regional temperature-change potentials for near-term climate forcings *Atmos. Chem. Phys.* **13** 2471–85
- Collins W J, Sitch S and Boucher O 2010 How vegetation impacts affect climate metrics for ozone precursors *J. Geophys. Res. Atmos.* **115** D23308
- Comyn-Platt E *et al* 2018 Permafrost and natural methane feedbacks limit emission budgets to  $1.5$  or  $2.0^\circ\text{C}$  of warming *Nat. Geosci.* in preparation
- Cox P and Jeffery H 2010 Methane radiative forcing controls the allowable  $\text{CO}_2$  emissions for climate stabilization *Curr. Opin. Environ. Sustain.* **2** 404–8
- Etminan M, Myhre G, Highwood E and Shine K 2016 Radiative forcing of carbon dioxide, methane, and nitrous oxide: a significant revision of the methane radiative forcing *Geophys. Res. Lett.* **43** 12614–23

- Gasser T, Peters G P, Fuglestedt J S, Collins W J, Shindell D T and Ciais P 2017 Accounting for the climate-carbon feedback in emission metrics *Earth Syst. Dyn.* **8** 235–53
- Gernaat D, Calvin K, Lucas P, Luderer G, Otto S, Rao S, Streffer J and van Vuuren D 2015 Understanding the contribution of non-carbon dioxide gases in deep mitigation scenarios *Glob. Environ. Change* **33** 142–53
- Harper A *et al* 2018 Relative effectiveness of land-based mitigation strategies in stabilising climate change at 1.5 °C *Nat. Commun.* in preparation
- Huntingford C *et al* 2010 IMOGEN: an intermediate complexity model to evaluate terrestrial impacts of a changing climate *Geosci. Model Dev.* **3** 679–87
- Huntingford C, Cox P, Mercado L, Sitch S, Bellouin N, Boucher O and Gedney N 2011 Highly contrasting effects of different climate forcing agents on terrestrial ecosystem services *Phil. Trans. R. Soc. A Math. Phys. Eng. Sci.* **369** 2026–37
- Huntingford C *et al* 2017 Flexible parameter-sparse global temperature time profiles that stabilise at 1.5 and 2.0 °C *Earth Syst. Dyn.* **8** 617–26
- Joos F, Bruno M, Fink R, Siegenthaler U, Stocker T F and LeQuere C 1996 An efficient and accurate representation of complex oceanic and biospheric models of anthropogenic carbon uptake *Tellus B Chem. Phys. Meteorol.* **48** 397–417
- MacDougall A H, Eby M and Weaver A J 2013 If anthropogenic CO<sub>2</sub> emissions cease, will atmospheric CO<sub>2</sub> concentration continue to increase? *J. Clim.* **26** 9563–76
- Meinshausen M, Wigley T and Raper S 2011 Emulating atmosphere-ocean and carbon cycle models with a simpler model, MAGICC6 Part 2: applications *Atmos. Chem. Phys.* **11** 1457–71
- Millar R J, Fuglestedt J S, Friedlingstein P, Rogelj J, Grubb M J, Matthews H D, Skeie R B, Forster P M, Frame D J and Allen A R 2017 Emission budgets and pathways consistent with limiting warming to 1.5 °C *Nat. Geosci.* **10** 741
- Morice C P, Kennedy J J, Rayner N A and Jones P D 2012 Quantifying uncertainties in global and regional temperature change using an ensemble of observational estimates: The HadCRUT4 data set *J. Geophys. Res. Atmos.* **117** D08101
- Myhre G, Highwood E J, Shine K P and Stordal F 1998 New estimates of radiative forcing due to well mixed greenhouse gases *Geophys. Res. Lett.* **25** 2715–8
- Myhre G *et al* 2013 Climate change 2013: the physical science basis *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed Stocker T *et al* (Cambridge: Cambridge University Press) pp 659–740
- O'Neill B *et al* 2017 The roads ahead: Narratives for shared socioeconomic pathways describing world futures in the 21st century *Glob. Environ. Change* **42** 169–80
- Pachauri R K *et al* 2014 Climate change 2014: synthesis report *Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (Geneva: IPCC)
- Riahi K *et al* 2017 The shared socioeconomic pathways and their energy, land use, and greenhouse gas emissions implications: an overview *Glob. Environ. Change* **42** 153–68
- Rogelj J, Luderer G, Pietzcker R, Krieglner E, Schaeffer M, Krey V and Riahi K 2015a Energy system transformations for limiting end-of-century warming to below 1.5 °C *Nat. Clim. Change* **5** 519
- Rogelj J, Meinshausen M, Schaeffer M, Knutti R and Riahi K 2015b Impact of short-lived non-CO<sub>2</sub> mitigation on carbon budgets for stabilizing global warming *Environ. Res. Lett.* **10** 075001
- Shindell D T, Fuglestedt J S and Collins W J 2017 The social cost of methane: theory and applications *Faraday Discuss.* **200** 429–51
- Sitch S, Cox P M, Collins W J and Huntingford C 2007 Indirect radiative forcing of climate change through ozone effects on the land-carbon sink *Nature* **448** 791–U4
- Sitch S *et al* 2008 Evaluation of the terrestrial carbon cycle, future plant geography and climate-carbon cycle feedbacks using five dynamic global vegetation models (DGVMs) *Glob. Change Biol.* **14** 2015–39
- Smith P 2016 Soil carbon sequestration and biochar as negative emission technologies *Glob. Change Biol.* **22** 1315–24
- Stohl A *et al* 2015 Evaluating the climate and air quality impacts of short-lived pollutants *Atmos. Chem. Phys.* **15** 10529–66
- Taylor K E, Balaji V, Hankin S, Juckes M, Lawrence B and Pascoe S 2013 CMIP5 data reference syntax (DRS) and controlled vocabularies
- van Vuuren D *et al* 2017 Energy, land-use and greenhouse gas emissions trajectories under a green growth paradigm *Glob. Environ. Change* **42** 237–50
- West J J, Fiore A M and Horowitz L W 2012 Scenarios of methane emission reductions to 2030: abatement costs and co-benefits to ozone air quality and human mortality *Clim. Change* **114** 441–61